a 2006 0264 1 of 1

The invention refers to medicine, namely to neurology and rehabilitation neurology.

The method of conservative treatment of diskal hernia consists in that it is carried out the curative gymnastics with the positional correction of diskogenic disorders, then the classical curative segmental massage paravertebrally by the sparing method. There are also administered during one month the preparations Lymphomyosot, Gallium-Heel, China-Homaccord for 10...15 drops 3 times a day per os and Spascupreel for 1 suppository 2 times a day during 10 days per rectum, and subcutaneously it is administered Discus Compositum for 2,2...3,3 ml 10 injections in a day, Colocynthis-Homaccord for 1,1...2,2 ml 5...10 injections in 2 days, Traumeli S for 2,2...3,2 ml 10 injections in a day, Coenzyme Compozitum for 2,2 ml 5 injections 2 times a week, Ubichinon Compositum for 2,2 ml 5 injections 2 times a week, at the same time the first 3 days the preparations are administered in the region of biologically active points G14, 10, 11, P7, MC6, TR5, E36, VB34, 39, RP6, C3-7, V62, then daily in the region of 6...8 biologically active points: IG3, G111, V11, VB34, IG3, IG7, IG8, V58, TR10, R8, T3-5, T14, V23-28, 31-34, 60. Within 20 days from the beginning of treatment it is administered Zeel P and Placenta Compositum for 2,2...3,3 ml 5...10 injections once in 10 days subcutaneously in the region of biologically active points: T3-5, 14, V23-28, 31, 34, 60, at the same time into each biologically active point it is administered only one mentioned medicinal preparation.

Claims: 1