

The invention refers to medicine, namely to neurology, pediatry and is meant for treatment of neurotic enuresis to children.

The method, according to the invention, consists in that the patient is administered *per os Berberis-Homaccord* and *Sabal-Homaccord* during 3 months 3 times a day 3...7 drops, *Reneel* during 1...2 months 1 tablet 3 times a day, *Nervoheel* during 2...3 months 1 tablet 3 times a day and subcutaneously *Cerebrum Compositum N* 2 times a week 1,1...2,2 ml, *Tonico-Injeel* 0,7...1,1 ml, *Solidago Compositum S* and *Traumeel S* 1,1...2,2 ml once a week during 5...10 weeks in the region of biologically active points E 36; IG 3; G 14; C 5; VB 39; TR 5, 10; P 10; V 21, 22, 28, 31, 40, 62; R 2, 7; F 2, 3, 8; VC 3, 4, 6; VG 2, 4, at the same time in each biologically active point it is administered only one indicated medicinal preparation, each preparation is administered in 3...5 points. It is carried out the general relaxing massage, it is limited the intake of liquids after 18.00 – 19.00 o'clock, it is awaken the child to 1.00 – 2.00 o'clock in the morning for miction, there are used sound signals for formation of the conditioned reflex for miction, it is kept the salt-free diet. The treatment course is repeated 2...3 times in 2...4 months.

Claims: 1